

Unsuccessful Infertility Treatment and Psychological Outcomes: A Prospective Cohort Study

Mehrnoosh Kakavand¹, Niloofar Gholami², Kobra Hosseini^{*1} , Atefeh Zandifar³, Haniyeh Rashidi¹, Fatemeh Rahimikia⁴, Hadith Rastad²

1. Clinical Research and Development Center of the Kamali Hospital, Alborz University of Medical Sciences, Karaj, Iran.

2. Cardiovascular Research Centre, Alborz University of Medical Sciences, Karaj, Iran.

3. Imam Ali Hospital, Alborz University of Medical Sciences, Karaj, Iran.

4. Iranian Center of Neurological Research, Neuroscience Institute, Tehran University of Medical Sciences.

ABSTRACT

Background and Aim: To quantitatively examine the association between the negative outcome of infertility treatment and psychological disorders in infertile women and to determine any difference in their post-treatment mental health compared to those who had a successful cycle.

Methods: This study included sixty-one infertile women who underwent infertility treatment between December 2022 and July 2023. Their mental health status was assessed with three self-reported measurement tools before the treatment and again after identifying the pregnancy test result.

Results: Women who had a positive test result were recruited into the successful group (n= 13), and those with a negative test were included in the unsuccessful group (n= 48). Women who had an unsuccessful treatment showed a significant increase in their levels of depression, anxiety, stress, hopelessness, and grief (P-value < 0.001). In comparison between women with successful and unsuccessful treatment, infertile women in the unsuccessful group showed higher levels of depression, anxiety, stress, hopelessness, and grief after the treatment (P-value < 0.001).

Conclusion: An unsuccessful treatment may potentially increase infertile women's psychological disorders. This suggests that addressing the psychological aspect alongside infertility treatments for infertile women may help achieve a comprehensive treatment approach.

ARTICLE INFO

Date Submitted: 13 January 2025

Date Accepted: 05 March 2025

KEYWORDS

Mental health; Infertility; IVF/ICSI;
Psychological disorders

*CORRESPONDING AUTHOR

Kobra Hosseini

Email: k.hosseini@abzums.ac.ir

 0009-0004-9261-126X

INTRODUCTION

Infertility is a significant concern in the field of reproductive health (1, 2). Despite recent progress in Assisted Reproductive Technology (ART), the global infertility rate continues to rise annually (3), and the rate of successful treatment is still low (4).

Psychological factors, including high levels of stress, anxiety, and depression, are among the identified predictors of treatment failure in infertile women (5-8).

A large percentage of infertile women experience one or more unsuccessful treatment processes. A qualitative study in 2019 reported that infertile individuals encountered stressors following unsuccessful treatment cycles. They described feeling exhausted from the treatment process, frustrated, fearful of failure of subsequent treatment, and stressed due to their relationship with their spouses (9).

However, few studies have quantitatively examined the impact of these experiences on their mental health, mainly addressing one aspect of psychological issues (10-12).



Please Cite This Paper As:

Kakavand M, Gholami N, Hosseini K, Zandifar A, Rashidi H, Rahimikia F, Rastad H. Unsuccessful Infertility Treatment and Psychological Outcomes: A Prospective Cohort Study. Stud Res Transl Med. 2025;7:e47246.

Open Access Policy: This article is distributed under the terms of the Creative Commons Attribution-Noncommercial 4.0 International License (<https://creativecommons.org/licenses/by-nc/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source and is not used for commercial purposes.



If an unsuccessful result leads to psychological problems, it could affect the patient's lifestyle and subsequent infertility treatments (13).

Given the importance of the issue and the absence of comprehensive evidence, this study aimed to examine the effect of unsuccessful infertility treatment on the levels of stress, anxiety, depression, feelings of hopelessness, and grief in infertile women.

MATERIALS and METHODS

This prospective cohort study was conducted at the Ibn Sina Infertility Treatment Center in Kamali Hospital, a tertiary care center specializing in reproductive medicine.

Study Design and Participants

After obtaining approval from the Ethics Committee, this cohort study recruited women who had undergone IVF/ICSI infertility treatment at the Ibn Sina Infertility Treatment Center from December 2022 to July 2023. The inclusion criteria for the study were as follows:

Women with a history of primary or secondary infertility seeking IVF (In Vitro Fertilization) treatment for the first time. No prior history of undergoing IVF or other assisted reproductive treatments (e.g., microinjection).

No specific etiology for infertility was required; meaning women with various causes such as male-factor, female-factor, or unexplained infertility were included.

Additionally, participants had to meet the following criteria: Be a candidate for infertility treatment during the recruitment period. Be older than 18 years. Provide informed consent to participate in the study.

The exclusion criteria were as follows

History of neurological or psychiatric problems or use of psychotropic medications.

Any problems in various aspects of the patient's life during the interval between two phases of data collection, such as the death of relatives, new challenges in personal life, or marital relationships.

Women who met the inclusion and exclusion criteria received counseling about study enrollment through both oral and written information and then signed a written informed consent after counseling. A comprehensive checklist covering sociodemographic information, medical information, and reproductive history was obtained from each participant via structured interview at the first visit to the clinic. The checklist consisted of the participant's age, education levels, employment and economic status, duration of marriage and infertility, history of puncture, number of live children and pregnancies, and the demographic information of their spouses (i.e., age, education, and

occupation). Additionally, the participants filled out the data collection tools in the interview room located at the IVF unit of the clinic. Researchers clarified questions that were not fully understood. The questionnaires were completed by participants at the beginning of the treatment and again after identifying the pregnancy test results. Every woman underwent Human Chorionic Gonadotropin (HCG) pregnancy blood testing 14 days after Intra- cytoplasmic sperm injection (ICSI). A positive pregnancy was determined by a serum HCG level exceeding 10 IU/L.

Measurement Tools

Beck Hopelessness Scale (BHS)

The Beck Hopelessness Scale (BHS) is a 20-question questionnaire developed by Beck et al. to measure hopelessness (14). Specific questions earn points for "yes" or "no" responses, with total scores ranging from 0 to 20. Higher scores indicate greater levels of hopelessness. The Persian version of BHS, translated and validated by Kaviani H et al., was utilized (15).

Depression Anxiety Stress Scales (DASS-21)

The DASS-21 is a shorter version of DASS-42, measuring Depression, Anxiety, and Stress. Scores range from 0 to 4, with each subscale score multiplied by two for comparability (16). Higher scores indicate higher levels of each condition. The Persian DASS-21 version was used in this study to evaluate participants' mental health (17).

Grief Experience Questionnaire (GEQ)

Barrett and Scott created the Grief Experience Questionnaire (GEQ) in 1989 with 55 items using a 5-point Likert scale. It evaluates various grief aspects like guilt, responsibility, and reactions to death (18). Mehdi-Poor et al. adapted the questionnaire for an Iranian study, resulting in a 34-item version with seven subscales. Responses are on a 5-point Likert scale, generating subscale scores from 5 to 25. Total scores of 34-68, 68-102, and over 102 indicate low, moderate, and high grief levels (19).

Statistical Assessment

Continuous variables are presented as mean $\pm$ SD or median (interquartile range). Categorical variables are presented as percentages. Chi-squared or Fisher's exact test was used to compare categorical variables. The t-test was used for normally distributed continuous variables, and the Mann-Whitney U test was used for non-parametric variables. ANCOVA was used to adjust for covariates to assess the effects of infertility treatment on psychological factors. A P-value of < 0.05 was considered statistically significant for all analyses. Analysis was performed using IBM SPSS v22.

RESULTS

A total of 61 women were included in the current study, and all completed two weeks of follow-up (loss to follow-up: 0%). The median ages of included participants and their spouses were 34 (28, 38) and 37 (33, 42), respectively. Participants were categorized into two groups based on the outcome of the treatment: successful (women who achieved pregnancy) and unsuccessful (women who did not). Of the participants, 48 were in the unsuccessful group, while 13 were in the successful group.

As presented in Table 1, the baseline characteristics of participants achieving successful pregnancy were comparable to those with unsuccessful ICSI (P -value > 0.05).

The comparison of DASS-21 after infertility treatment demonstrated that patients in the unsuccessful group exhibited higher scores of depression, anxiety, and stress compared to the successful group (mean difference (95% CI): 20.00 (14.00, 26.00), 16.00 (10.00, 22.00), and 20.00 (14.00, 22.00), respectively). Furthermore, women with unsuccessful pregnancies experienced higher levels of hopelessness (mean difference (95% CI): 14 (12.27, 15.00)) and grief (mean difference (95% CI): 37.36 (27, 57)) after the treatment in comparison with those who achieved pregnancy (Table 2). Changes in the psychological status of all patients are summarized in Figures 1-2.

After adjusting for confounding variables, including the baseline psychological scores, age, occupation, and education levels of women and their spouses, number of abortions, live children, and stillbirth, duration of infertility and marriage, history of puncture, and their socioeconomic status, mean differences in all three subscales of DASS-21, Beck hopelessness, and GEQ-34 scores remained statistically significant (Table 2).

The evaluation of the median of three subscales of DASS-21, BHS, and GEQ-34 scores before and after the treatment in the successful and unsuccessful groups revealed the following: In the unsuccessful group, the median depression score increased from 5.50 to 27, anxiety from 5 to 21, stress from 11 to 26.85, BHS from 2.72 to 15.72, and GEQ-34 from 61.84 to 83.39. Conversely, for women in the successful group, the median depression score remained unchanged, anxiety and stress decreased from 8 to 6 and 12 to 8, respectively, and GEQ-34 decreased from 48.35 to 46.00 (Table 2).

Not only were the mean scores of DASS-21, BHS, and GEQ-34 significantly higher in women with unsuccessful ICSI, but all the aforementioned expected scores also had a predominantly increasing trend in non-pregnant patients. In contrast, they had a decreasing trend in pregnant patients (Figures 1-2).




FIGURE 1. A, B, C: Changes in levels of depression, anxiety, and stress according to DASS-21 for each participant.




FIGURE 2. A: Changes in levels of hopelessness according to BHS for each participant. B: Changes in levels of grief (all items) according to GEQ-34 for each participant.

Table 1. Characteristics

Descriptive characteristics		Successful group N = 13	Unsuccessful group N = 48	Total N = 61	P-value
Age (years) Median (IQR)		33 (27, 38)	34.5 (28, 39)	34 (28, 38)	0.432
Education n (%)	Below diploma	23.1 (3)	29.2 (14)	27.9 (17)	0.791
	Highschool diploma	30.8 (4)	35.4 (17)	34.4 (21)	
	Academic	46.2 (6)	35.4 (17)	37.7 (23)	
Employment status n (%)	House Wife	84.6 (11)	85.4 (41)	85.2 (52)	0.780
	Employee	15.4 (2)	10.4 (5)	11.5 (7)	
	Self-Employed	0.0 (0)	4.2 (2)	3.3 (2)	
Socio-economic status n (%)	Very low	7.7 (1)	0.0 (0)	1.6 (1)	0.367
	Low	0.0 (0)	6.2 (3)	4.9 (3)	
	Moderate	92.3 (12)	81.2 (39)	83.6 (51)	
	High	0.0 (0)	8.3 (4)	6.6 (4)	
	Very high	0.0 (0)	4.2 (2)	3.3 (2)	
Infertility n (%)	Primary	69.2 (9)	69.8 (30)	69.6 (39)	> 0.99
	Secondary	30.8 (4)	30.2 (13)	30.4 (17)	
Infertility Duration Median (IQR)		3 (2,4)	4 (2,6)	4 (2,6)	0.295
Years of Marriage Median (IQR)		6 (3.75, 10.5)	7 (3.75, 10.25)	6.5 (3.75, 10.25)	0.952
History of puncture n (%)	Yes	30.8 (4)	25.0 (12)	26.2 (16)	0.728
	No	69.2 (9)	75.0 (36)	73.8 (45)	
Number of Pregnancy Median (IQR)		1 (0, 1.25)	1 (0, 1.5)	1 (0, 1.5)	0.811
Live Children Median (IQR)		0 (0, 0.25)	0 (0, 1)	0 (0, 1)	0.708
Abortion Median (IQR)		0.5 (0, 1)	0 (0, 0.75)	0 (0, 1)	0.281
Still Birth Median (IQR)		0 (0, 0)	0 (0, 0)	0 (0, 0)	0.359
Characteristics of spouses					
Age (years) mean $\pm$ SD		35 (32, 37)	39 (33, 45.25)	37 (33, 42)	0.069
Education n (%)	Below diploma	38.5 (5)	37.5 (18)	37.7 (23)	> 0.99
	Highschool diploma	30.8 (4)	35.5 (17)	34.4 (21)	
	Academic	30.8 (4)	27.1 (13)	27.9 (17)	
Employment status n (%)	Employee	30.8 (4)	50.0 (24)	45.9 (28)	0.357
	Self-Employed	69.2 (9)	50.0 (24)	54.1 (33)	

IQR: Interquartile Range; SD: Standard Deviation

Table 2. Physiological status

Instrument		Successful Median (IQR)	Unsuccessful Median (IQR)	P-value (crude)	P-value (Adjusted model) *
Depression	Pre	6.00 (2.00, 14.00)	5.50 (2.00, 10.50)	0.582	
	Post	6.00 (2.00, 8.00)	27.00 (16.00, 32.50)	< 0.001	0.001
	Post-Pre (MD (95%CI))	-2.85 (-7.47, -7.47)	18.00 (15.00, 15.00)		
	(P-Value)	0.203	< 0.001		
Stress	Pre	12 (7.00, 19.10)	11 (4.00, 18.00)	0.627	
	Post	8.00 (4.00, 10.00)	26.85 (18.00, 34.00)	< 0.001	0.001
	Post-Pre (MD (95%CI))	-4.16 (-8.67, -8.67)	15.00 (12.00, 12.00)		
	(P-Value)	0.067	< 0.001		
Anxiety	Pre	8.00 (2.00, 13.00)	5.00 (2.00, 10.50)	0.456	
	Post	6.00 (2.00, 6.00)	21.00 (13.60, 28.50)	< 0.001	0.019
	Post-Pre (MD (95%CI))	-3.09 (-6.74, -6.74)	13.00 (11.00, 11.00)		
	(P-Value)	0.089	< 0.001		
Hopelessness	Pre	1.00 (1.00, 4.05)	2.72 (1.06, 5.00)	0.338	
	Post	2.00 (1.00, 3.00)	15.72 (14.00, 17.00)	< 0.001	<0.001
	Post-Pre (MD (95%CI))	-0.53 (-5.00, -5.00)	12.83 (11.50, 11.50)		
	(P-Value)	0.552	< 0.001		
Guilt	Pre	13.00 (10.00, 14.00)	12.00 (9.75, 15.00)	0.965	
	Post	11.00 (10.33, 13.00)	19.22 (14.75, 28.25)	< 0.001	0.053
	Post-Pre (MD (95%CI))	-1.00 (-3.34, -3.34)	6.34 (4.61, 4.61)		
	(P-Value)	0.683	< 0.001		
Search for explanation	Pre	9.00 (8.00, 11.00)	13.00 (10.75, 17.00)	0.023	
	Post	9.00 (7.00, 9.00)	17.00 (14.75, 22.25)	< 0.001	0.040
	Post-Pre (MD (95%CI))	-1.50 (-3.67, -3.67)	4.34 (2.88, 2.88)		
	(P-Value)	0.099	< 0.001		
Somatic reaction	Pre	6.00 (5.00, 10.00)	9.48 (7.00, 12.00)	0.08	
	Post	7.00 (6.00, 10.00)	12.79 (10.00, 18.00)	< 0.001	0.036
	Post-Pre (MD (95%CI))	-0.50 (-3.72, -3.72)	3.72 (2.50, 2.50)		
	(P-Value)	0.798	< 0.001		
Loss of social support	Pre	6.00 (4.00, 7.00)	7.09 (5.00, 9.00)	0.13	
	Post	5.00 (4.00, 6.00)	10.00 (8.00, 14.00)	< 0.001	0.052
	Post-Pre (MD (95%CI))	-1.35 (-2.50, -2.50)	3.41 (2.50, 2.50)		
	(P-Value)	0.065	< 0.001		
Stigmatization	Pre	4.00 (3.00, 4.00)	4.71 (3.00, 7.00)	0.036	
	Post	3.00 (3.00, 4.00)	7.00 (6.00, 10.25)	< 0.001	0.293
	Post-Pre (MD (95%CI))	0.11 (-2.00, -2.00)	3.50 (2.00, 2.00)		
	(P-Value)	0.888	< 0.001		

Instrument		Successful Median (IQR)	Unsuccessful Median (IQR)	P-value (crude)	P-value (Adjusted model) *
Shame	Pre	6.00 (5.00, 7.00)	7.00 (4.75, 8.25)	0.594	
	Post	4.00 (4.00, 7.00)	9.00 (8.00, 14.00)	< 0.001	0.159
	Post-Pre (MD (95%CI))	-1.00 (-3.00, -3.00)	3.00 (2.25, 2.25)		
	(P-Value)	0.138	< 0.001		
Judgment	Pre	5.00 (5.00, 6.00)	6.56 (4.00, 7.00)	0.375	
	Post	5.00 (4.00, 6.00)	10.00 (8.00, 14.25)	< 0.001	0.202
	Post-Pre (MD (95%CI))	-1.00 (-3.50, -3.50)	4.00 (2.50, 2.50)		
	(P-Value)	0.598	< 0.001		
Total GEQ-34	Pre	48.35 (44.58, 55.00)	60.84 (48.75, 73.25)	0.043	
	Post	46.00 (42.00, 50.00)	83.39 (71.91, 118.35)	< 0.001	0.045
	Post-Pre (MD (95%CI))	-5.65 (-13.35, -13.35)	21.14 (16.16, 16.16)		
	(P-Value)	0.078	< 0.001		

*For the P-value (adjusted model), we used ANCOVA to adjust covariates, including the age of women, education of women, women's jobs, number of pregnancies, live children, abortions, stillbirths, infertility type, and infertility duration, years of marriage, history of puncture, successful puncture, socioeconomic status, spouses' age, education, and jobs, and patients' psychological scores before treatment. IQR: Interquartile Range; MD: Mean Difference; Pre: before treatment; Post: after treatment; 95% CI: 95% Confidence Interval; GEQ-34: Grief Experience Questionnaire 34.

DISCUSSION

The present cohort study aimed to identify if unsuccessful infertility treatment outcomes could predict psychological distress in infertile women. Our findings revealed a significant association between unsuccessful treatment and increased levels of depression, anxiety, stress, feelings of hopelessness, and grief in infertile individuals, considering pre-ICSI mental health status.

These findings were similar to Pasch et al. and Verhaak et al., both of whom observed a significant increase in levels of depression (from 12.39 to 15.85 and 1.3 to 3.9, respectively) and anxiety (from 41.41 to 43.33 and 36.9 to 39.8, respectively) in their participants following a failed IVF cycle (10, 12). Additionally, Pasch et al. argued that levels of depression following unsuccessful treatment were comparable to those experienced by parents who had lost a child and even higher than those reported for cancer patients undergoing treatment (10). Although in these two studies, unsuccessful treatment led to a significant increase in psychological problems in infertile women, the initial increase in psychological problems of unsuccessful women in our study was greater than theirs. In the context of feelings of hopelessness and grief, the median score of BHS in the unsuccessful group after the failed treatment was even higher than breast cancer patients and vulnerable prisoners (mean: 13.3, 10.13, respectively) (14, 15). These women also showed a higher median score of grief than Iranian

individuals who experienced the death of relatives during the COVID-19 pandemic (mean: 80.27) (16).

The higher increases in psychological scores following a failed infertility treatment in our study compared to similar studies in other communities may stem from cultural distinctions among different populations. Iran, as a Middle Eastern country, places a strong emphasis on a woman's social standing, dignity, and self-worth concerning her ability to conceive (32). A recent study showed the high level of anxiety disorders, which are mainly linked to economic hardships in Iran (17). Given the high cost of infertility treatment, the stress and fear of a negative outcome and the cost of the next treatment could explain the significant psychological pressure on infertile individuals in such countries.

The increased psychological problems following unsuccessful infertility treatments may negatively affect the motivation for subsequent treatments and the overall lifestyle of infertile individuals (18). In this regard, Verhaak et al. showed that unsuccessful treatment not only increased levels of psychological distress in infertile women but also served as a less favorable starting point for the next treatment cycle (13). Also, based on recent evidence, these induced mental health problems could negatively impact the results of subsequent treatment (19-21). The mechanism by which mental disorders can lead to unsuccessful infertility treatment remains unclear. However, it has been suggested that mental disorders may influence the function of endocrine glands, the

immune system, and oocyte retrieval outcomes, all of which are associated with reduced fertility and decreased chances of successful treatment (20, 22).

It is established that a high level of depression shortly after a stressful event is a significant risk factor for the development of a depressive disorder (23). Although limited longitudinal studies with constraints such as high drop-out rates showed a decreasing trend over time for levels of psychological disorders following a failed treatment, a significant percentage of women, nearly twenty percent, remain depressed even six months after the last failure (24).

On the other hand, while the levels of psychological issues in our study decreased after treatment in the successful group, a positive result did not alleviate psychological disorders in all infertile women, as some still experienced significant levels of depression, stress, anxiety, as well as feelings of hopelessness and grief (Table 2). This is in line with earlier studies and is supported by research suggesting that the fear of miscarriage may persist even after receiving a positive pregnancy, which could explain the persistence of psychological issues (13, 25).

Healthcare providers need to understand that treatment procedures and their failure can exacerbate the trauma experienced by infertile women. Therefore, taking an approach that encompasses the emotional and psychological well-being of patients, such as counseling during infertility treatment, may be helpful as part of the treatment procedure for women undergoing infertility treatment, especially in societies where motherhood holds significant social influence for women.

CONCLUSION

Infertility, often perceived through the narrow lens of biomedical failure, exacts a far deeper toll—one etched into the psyche of those who endure its uncertainties. This prospective cohort study elucidates the profound psychological ramifications of unsuccessful IVF/ICSI treatment, revealing a stark dichotomy: while successful conception alleviates distress, treatment failure precipitates a surge in depression, anxiety, stress, hopelessness, and grief. The magnitude of this deterioration is not merely statistically significant but clinically alarming, with post-treatment psychological scores in the unsuccessful cohort eclipsing those observed in populations grappling with severe illness or bereavement.

The implications extend beyond transient emotional distress. In cultures where fertility is inextricably linked to identity and social capital—such as Iran, the setting of this study—the psychological burden is compounded by existential and

societal pressures. The data suggest that infertility treatment, when unsuccessful, does not merely fail to achieve pregnancy; it risks destabilizing mental health in ways that may undermine future treatment adherence, marital dynamics, and overall well-being. Notably, even "success" does not universally confer psychological relief, as residual anxiety persists in some women, likely fueled by the fragility of early pregnancy after prolonged infertility.

These findings demand a paradigm shift in reproductive care. The current model, which prioritizes biochemical success over holistic well-being, is insufficient. Integrating evidence-based mental health support—such as preemptive counseling, mindfulness-based stress reduction, or structured grief interventions—into standard infertility protocols could mitigate the collateral damage of treatment failure. Moreover, fostering cross-disciplinary collaboration between reproductive endocrinologists and mental health professionals is imperative to address the bidirectional relationship between psychological distress and treatment outcomes.

Future research should investigate longitudinal mental health trajectories post-treatment and the efficacy of targeted psychosocial interventions. For now, this study serves as a clarion call: infertility is not solely a disorder of the body but a crisis of the self. Medicine must respond with equal rigor to both dimensions.

ACKNOWLEDGEMENTS

Researchers thanked Clinical Research Development Units at Rajaie and Kamali Hospitals, Alborz University, and Dr. Razieh Lotfi for manuscript review support.

CONFLICT OF INTEREST

The authors declare no competing interests.

FUNDING

There were no grants or financial support obtained for this study.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the corresponding author upon reasonable request.

REFERENCES

1. Kaur A. Role of Ayurvedic herbs for healthy pregnancy. *International Journal of Research in Engineering, Science and Management*. 2022;5(12):16-7.

2. Kawamura K, Sato Y. Effectiveness of supplement ingredients on infertility treatment in advanced aged women. *Journal of Obstetrics and Gynaecology Research*. 2023.
3. Naeem A, Khan MZ, Khan A, Abbas SH, Khan S. Amniotic stem cells as a source of regenerative medicine to treat female infertility. *Human Cell*. 2023;36(1):15-25.
4. Peng M, Zhang J, Zeng T, Hu X, Min Z. Stress, anxiety, and depression in infertile couples are not associated with a first IVF or ICSI treatment outcome. *BMC Pregnancy and Childbirth*. 2021;21(1):1-8.
5. Lynch CD, Sundaram R, Maisog JM, Sweeney AM, Buck Louis GM. Preconception stress increases the risk of infertility: results from a couple-based prospective cohort study—the LIFE study. *Human Reproduction*. 2014;29(5):1067-75.
6. Sominsky L, Sobinoff AP, Jobling MS, Pye V, McLaughlin EA, Hodgson DM. Linking stress and infertility: a novel role for ghrelin. *Endocrine Reviews*. 2017;38(5):432-67.
7. Smeenk JM, Verhaak CM, Vingerhoets AJ, Sweep CG, Merkus JM, Willemsen SJ, et al. The effect of anxiety and depression on the outcome of in-vitro fertilization. *Human Reproduction*. 2001;16(7):1420-3.
8. Klonoff-Cohen H, Chu E, Natarajan L, Sieber W. A prospective study of stress among women undergoing in vitro fertilization or gamete intrafallopian transfer. *Fertility and Sterility*. 2001;76(4):675-87.
9. Zagami SE, Latifnejad Roudsari R, Janghorban R, Mousavi Bazaz SM, Amirian M, Allan HT. A qualitative study of the experiences of Iranian infertile couples after unsuccessful assisted reproductive technologies. *International Journal of Women's Health and Reproduction Sciences*. 2019;7(3):331-8.
10. Pasch LA, Gregorich SE, Katz PK, Millstein SG, Nachtigall RD, Bleil ME, et al. Psychological distress and in vitro fertilization outcome. *Fertility and Sterility*. 2012;98(2):459-64.
11. Cheung CS, Chan CH, Ng EH. A prospective observational study on the stress levels at the time of embryo transfer and pregnancy testing following in vitro fertilisation treatment: a comparison between women with different treatment outcomes. *BJOG: An International Journal of Obstetrics & Gynaecology*. 2019;126(2):271-9.
12. Verhaak CM, Smeenk JM, Evers AW, Kremer JA, Kraaijmaat FW, Braat DD. Long-term psychological adjustment to IVF/ICSI treatment in women. *Human Reproduction*. 2007;22(1):305-8.
13. Verhaak CM, Smeenk JM, van Minnen A, Kremer JA, Kraaijmaat FW. Stress and marital satisfaction among women before and after their first cycle of in vitro fertilization and intracytoplasmic sperm injection. *Fertility and Sterility*. 2001;76(3):525-31.
14. Palmer EJ, Connelly R. Depression, hopelessness and suicide ideation among vulnerable prisoners. *Criminal Behaviour and Mental Health*. 2005;15(3):164-70.
15. Bener A, Alsulaiman R, Doodson L, El Ayoubi HR. Depression, hopelessness and social support among breast cancer patients: in highly endogamous population. *Asian Pacific Journal of Cancer Prevention*. 2017;18(7):1889.
16. Shahini N, Ebrahimzadeh A, Shahini N. Grief experience after deaths: Comparison of COVID-19 and non-COVID-19 causes. *Journal of Patient Experience*. 2022;9:23743735221089697.
17. Moradinazar M, Najafi F, Jalilian F, Pasdar Y, Hamzeh B, Shakiba E, et al. Epidemiological status of depressive disorders in the Middle East and North Africa from 1990 to 2019. *Health Promotion Perspectives*. 2022;12(3):301.
18. Vikström J, Josefsson A, Bladh M, Sydsjö G. Mental health in women 20–23 years after IVF treatment: a Swedish cross-sectional study. *BMJ Open*. 2015;5(10):e009426.
19. Aimagambetova G, Terzic S, Laganà AS, Bapayeva G, Ukybassova T, Aldiyarova A, et al. The effect of psychological distress on IVF outcomes: Reality or speculations? *PLoS One*. 2020;15(12):e0242024.
20. Ebbesen SM, Zachariae R, Mehlsen MY, Thomsen D, Højgaard A, Ottosen L, et al. Stressful life events are associated with poor in-vitro fertilization (IVF) outcome: a prospective study. *Human Reproduction*. 2009;24(9):2173-82.
21. Thiering P, Beaupaire J, Jones M, Saunders D, Tennant C. Mood state as a predictor of treatment outcome after in vitro fertilization/embryo transfer

- technology (IVF/ET). *Journal of Psychosomatic Research*. 1993;37(5):481-91.
22. Szkodziak F, Krzyżanowski J, Szkodziak P. Psychological aspects of infertility: A systematic review. *Journal of International Medical Research*. 2020;48(6):0300060520932403.
 23. Horwath E, Johnson J, Klerman GL, Weissman MM. What are the public health implications of subclinical depressive symptoms? *Psychiatric Quarterly*. 1994;65:323-37.
 24. Verhaak CM, Smeenk JM, Evers AW, Kremer JA, Kraaijmaat FW, Braat DD. A longitudinal, prospective study on emotional adjustment before, during, and after consecutive fertility treatment cycles. *Human Reproduction*. 2005;20(8):2253-60.
 25. Volgsten H, Skoog Svanberg A, Ekselius L, Lundkvist O, Sundström Poromaa I. Risk factors for psychiatric disorders in infertile women and men undergoing in vitro fertilization treatment. *Fertility and Sterility*. 2010;93(4):1088-96.