


Journal Pre-proof

Association of detectable C-peptide levels with glycemic control and chronic complications in individuals with type 1 diabetes mellitus: A systematic review and meta-analysis

Mahin Seifi Alan, Amirhossein Tayebi, Elmira Jafari Afshar, Sanaz Seifi Alan, Mahnaz Seifi Alan, Ramina Fazeli, Tooba Sohbatzade, Parham Samimisedeh, Hadith Rastad



PII: S1056-8727(24)00193-4

DOI: <https://doi.org/10.1016/j.jdiacomp.2024.108867>

Reference: JDC 108867

To appear in: *Journal of Diabetes and Its Complications*

Received date: 7 April 2024

Revised date: 22 August 2024

Accepted date: 14 September 2024

Please cite this article as: M.S. Alan, A. Tayebi, E.J. Afshar, et al., Association of detectable C-peptide levels with glycemic control and chronic complications in individuals with type 1 diabetes mellitus: A systematic review and meta-analysis, *Journal of Diabetes and Its Complications* (2024), <https://doi.org/10.1016/j.jdiacomp.2024.108867>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Title:**Association of detectable C-peptide levels with glycemic control and chronic complications in individuals with type 1 diabetes mellitus: A systematic review and meta-analysis****Running title:** Clinical outcomes of C-peptide in type 1 diabetes

Mahin Seifi Alan¹, Amirhossein Tayebi^{1&2}, Elmira Jafari Afshar¹, Sanaz Seifi Alan¹, Mahnaz Seifi Alan¹, Ramina Fazeli¹, Tooba Sohbatzade¹, Parham Samimisedeh¹, and Hadith Rastad^{1&2}.

1 Cardiovascular Research Center, Alborz University of Medical Sciences, Karaj, Iran;

2 Non-communicable Diseases Research Center, Alborz University of Medical Sciences, Karaj, Iran.

* Parham Samimisedeh and Hadith Rastad contributed equally as corresponding authors, Cardiovascular Research Center, Alborz University of Medical Sciences, Karaj, Iran; e-mail: parhamsamimimd@gmail.com & h.rastad91@gmail.com; Postal code: 3149779453; Tel: +9826 32563318.

Highlights

- We have summarized the available evidence on the clinical significance of detectable levels of C-peptide in T1DM.
- A systematic search and meta-analysis were performed using online databases.
- Individuals with T1DM in the detectable C-peptide group, compared with the undetectable C-peptide group, had lower mean HbA1c (- 0.08) and daily insulin dose (- 0.41) and showed lower odds for retinopathy (pooled crude odds ratios: 0.53) and nephropathy complications (0.62).
- Individuals with T1DM in the detectable C-peptide group may experience better clinical outcomes.

Journal Pre-proof

Manuscript word count: 3386

Abstract word count: 231

Author Disclosures: The authors declare no conflict of interest.

Acknowledgments: The researchers appreciate the Clinical Research Development Units of Kamali and Rajaei Hospitals in Alborz University of Medical Sciences.

Journal Pre-proof

Abstract:

Aims: Multiple studies have addressed the association between detectable levels of C-peptide and glycemic control, as well as the development of chronic complications of type 1 diabetes mellitus (T1DM), including both macrovascular and microvascular diseases. We aimed to summarize the available evidence on the clinical significance of detectable levels of C-peptide in T1DM.

Method: A systematic search was performed on online databases using the following key terms: T1DM, C-peptide, diabetes mellitus complications, and glycemic parameters. We pooled standardized mean difference (SMD) and odds ratios (OR).

Results: Of the 1,519 articles retrieved from the initial search, 38 (12 cohort and 26 cross-sectional studies) met our eligibility criteria. Individuals with T1DM in the detectable C-peptide group, compared with the undetectable C-peptide group, had lower mean HbA1c [pooled SMD (95% confidence interval (95% CI)): - 0.08 (- 0.13 to - 0.02), $I^2 = 0\%$, p.value: 0.005] and daily insulin dose [- 0.41 (- 0.65 to - 0.18), $I^2 = 83\%$, p.value < 0.001]. They also showed lower odds for retinopathy [pooled crude OR (95% CI): 0.53 (0.41 to 0.69), $I^2 = 65\%$, p.value < 0.001] and nephropathy complications [0.62 (0.55 to 0.70), $I^2 = 19\%$, p.value < 0.001]; however, the two groups were similar regarding neuropathy [0.92 (0.65 to 1.31), $I^2 = 0\%$, p.value: 0.31].

Conclusions: The available evidence suggests that individuals with T1DM in the detectable C-peptide group may experience better clinical outcomes.

Keywords: T1DM; C-peptide; glycemic control parameters; diabetes mellitus complications

1. Introduction

Type 1 diabetes mellitus (T1DM) is a condition characterized by the loss of β -cell mass in pancreatic islets caused by an autoimmune response ¹. The novel ultrasensitive method for measuring C-peptide levels allows for the detection of lower levels of this biomarker than previously possible in individuals with T1DM, even long after diagnosis ^{2,3}.

Many studies have reported detectable levels of C-peptide in individuals with T1DM years after their initial diagnosis, indicating the presence of endogenous insulin secretion by residual autoimmune-resistant β -cells ⁴⁻⁶. Previous studies have explored the risk factors and determinants associated with detectable C-peptide levels in T1DM. In this regard, some human leukocyte antigens (HLA) serotypes, including HLA DR3, DR4, and DQ8, along with early age at diagnosis, long-standing disease, higher HbA1c at the time of diagnosis, and male gender, are inversely associated with detectable C-peptide levels in individuals with T1DM ^{2,7}.

Increasing numbers of studies have assessed whether detectable C-peptide levels are correlated with lower glucose level fluctuations and better glycemic control in individuals with T1DM ^{8,9}. Additionally, some studies have addressed the association between detectable C-peptide levels and chronic complications of T1DM, including macrovascular and microvascular diseases ^{10,11}.

Nevertheless, current evidence regarding the clinical importance of detectable C-peptide levels in individuals with T1DM is scattered, making it difficult to utilize these data in clinical decision-making. Thus, as the first systematic review and meta-analysis in this area, our goal is to provide a clear and cohesive summary of the key findings from the existing evidence on the clinical significance of detectable C-peptide levels in individuals with T1DM.

2. Materials and Methods

We conducted a systematic review of the literature following a standardized methodology and reported our findings according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines. Since our study was a systematic review of existing literature, we were exempt from obtaining institutional ethics committee approval. We included all original studies assessing C-peptide levels in individuals with T1DM.

2.1. Search strategy

A systematic search was conducted on online databases, including PubMed, Embase, and Scopus, on July 16, 2024, using a combination of related keywords in three domains: T1DM, C-peptide, and diabetes complications (diabetic retinopathy, diabetic neuropathy, diabetic nephropathy, hypoglycemia, diabetic cardiomyopathy, foot ulcer, diabetic ketoacidosis (DKA), albuminuria, microalbuminuria, microaneurysms, and nerve conduction). We used the Boolean operator "OR" to connect key terms within each domain and the "AND" operator to connect the domains. In addition, we manually searched the first 100 pages of Google Scholar and the reference lists of relevant articles for any additional citations (Figure 1). We imported all citations from the retrieved documents into EndNote software (version X9.3.2, Captivate Analytics, California, USA) and removed duplicate articles.

2.2. Study selection

Two researchers (M.S1 & R.F) independently screened the titles, abstracts, and full texts of the imported articles to identify eligible studies. Any disagreements were resolved through discussion. To be included in our review study, studies had to report C-peptide levels in individuals with T1DM and meet the following criteria:

1. Individuals with T1DM
2. Presence of at least one measurement of C-peptide
3. Observational study design: cross-sectional studies, case-control, and cohort studies
4. Written in English

5. At least one of the following complications related to T1DM included in the studies: glycemic control, microvascular complications, or macrovascular complications.

Exclusion criteria

1. Animal studies, in vitro studies, and review articles
2. Not available in full text

2.3. Data extraction

We extracted the following data from the full text of the included articles into "Data extraction form" in Microsoft Excel (Version 2016, Microsoft Corp., Redmond, WA, USA): First author's name, publication year, country, study design, sample size, age, gender, BMI, time of diagnosis of T1DM, technique for assessing C-peptide, and the C-peptide levels.

2.4. Quality assessment

Two of our researchers (M.S1 & S.S) assessed the quality of the included studies independently using the Newcastle–Ottawa Scale (NOS) appraisal tool checklists adopted for each type of study design to assess the quality of the included studies. Our third researcher (H.R.) resolved any disagreements (Table 1 and 2).

2.5. Statistical analysis

We utilized either the random-effect or the fixed-effect models based on the heterogeneity size of the Standardized Mean Difference (SMD) or odds ratios (OR). The magnitude and significance of the heterogeneity were determined by I-squared statistics and Q-test, respectively. We used the random-effect model if the I-squared statistic was greater than 25%. We combined SMD and OR using inverse variance and Mantel-Haenszel methods, respectively. We computed the SMD for HbA1c and insulin dose, as well as the crude OR for retinopathy, nephropathy, and neuropathy. These calculations were performed for the comparison between the detectable and undetectable C-peptide groups among individuals with T1DM. Meta-analyses were conducted using the R Meta package in R Studio software (version 4.2.2).

3. Results

In the initial search, a total of 1,519 articles were identified. Following the removal of duplicates (238 articles) and those that did not meet the inclusion criteria based on title/abstract (1,127 articles) or full text (116 articles), 38 articles were considered eligible for this study. These 38 articles consisted of 12 cohort studies (mean/median follow-up duration of 0.5 years to 20 years) and 26 cross-sectional studies. Please refer to Figure 1 for a visual representation of this process.

Table 3 presents the characteristics of the included studies. Overall, the included studies investigated the associations between C-peptide status and glycemic control indicators (HbA1c (n = 27)^{2,4,5,7,8,11-32}, daily insulin dose (n = 20)^{2,4,8,14-20,22,23,26-29,31-34}, hypoglycemia events (n = 11)^{7,15,16,20,22,25,29,31,35-37}, DKA events (n = 4)^{16,17,31,38}, mean glucose (n = 6)^{8,14,18,25,26,32}, coefficient of variation (CV) (n = 7)^{8,18,21,25,26,28,39}, %time spent above (n = 5)^{8,14,25,26,32}/below (n = 4)^{8,14,21,26} and in range (n = 8)^{8,14,18,21,26,28,32,39}, glucose standard deviation (SD) (n = 2)^{18,25}, mean amplitude of glycemic excursions (MAGE) (n = 1)¹⁸, microvascular complications (overall (n = 2)^{7,13} or by type: nephropathy (n = 6)^{2,15,16,29,30,40}, retinopathy (n = 16)^{2,4,5,12,15-17,22,25,29,30,32,34,38,40-42}, albuminuria (n = 6)^{2,4,22,29,34,38}, neuropathy (n = 6)^{4,15,17,29,34,38}), and macrovascular complications (n = 5)^{2,4,13,15,43}, (overall, cardiovascular disease (CVD), cardiovascular event, or coronary artery disease (CAD)) in individuals with T1DM (Table 4).

Journal Pre-proof





Figure 1: PRISMA Flowchart for the selection process of studies

Included studies were published between 2008 and 2024^{2,5,13,14,16,19,20,22,25,26,28,33,35-37,41}. The number of participants varied from 19 individuals in Babaya et al.'s study³⁹ to 5,732 in Jeyam et al.'s study¹⁶, which were recruited using different inclusion criteria. The mean/median T1DM duration, reported by 33 studies, varies from 0 to 56 years and was lower than 20 years in 22 out of the 33 studies. In 21 studies, T1DM duration was compared between C-peptide groups, and 9 of them found a lower disease duration in individuals with T1DM in the detectable C-peptide group than in the undetectable group. Most research comparing gender (18 out of 22 studies) or BMI (18 out of 20 studies) between the two groups found no significant findings in this regard.


Studies utilized different types of C-peptide tests (reported by 36 studies), including random (rCP) (n = 7)^{2,5,16,24,25,35,40}, fasting (fCP) (n = 11)^{7,13,29-32,34,37-39,44}, and stimulated (n = 13)^{4,8,14,17-23,26,27,33}, i.e., postprandial C-peptide (PCP), mixed meal tolerance test (MMTT), oral glucose tolerance test (OGTT), and glucagon stimulation test (GST)). Additionally, five studies utilized both fasting and stimulated methods^{12,15,32,33,42}, and two studies used the urinary C-peptide creatinine ratio (UCPCR)^{28,43}. In addition, included studies categorized participants into two groups using different cut points for C-peptide (Table 4).

3.1. HbA1c

A total of 25 studies (6 cohorts^{12,14,16-19} and 19 cross-sectional^{2,4,5,7,8,13,15,20-29,31,44}) compared HbA1c between individuals with T1DM in the detectable and undetectable C-peptide groups; of those, 11 studies found a significant difference between detectable and undetectable C-peptide groups. All consistently showed that individuals with T1DM in detectable C-peptide group had a lower HbA1c^{2,4,7,12,15,16,18,20,21,23,24}. All three studies^{2,20,23} that accounted for different confounding variables, such as age at onset and duration of T1DM, reported a significant inverse association between C-peptide and HbA1c levels. A meta-analysis was conducted in three cohort studies^{13,17,38}, and nineteen cross-sectional studies^{2,4,5,7,8,13,15,20-29,31,44}. The pooled estimations of SMD indicated that HbA1c levels were lower in the detectable C-peptide group compared to the undetectable group, although this difference was not found to be statistically significant [pooled SMD (95% confidence interval (95% CI)): - 0.47 (- 1.05 to 0.10), $I^2 = 97\%$, p.value: 0.11]. However, in subgroup analysis by study design, this difference reached statistical significance in the meta-analysis of the cohort studies [pooled SMD (95% CI): - 0.08 (- 0.13 to - 0.02), $I^2 = 0\%$, p.value: 0.005] (Table 4, Figures 2 and 3a).



a



b

Figure 2: Pooled estimations of standardized mean difference (SMD) and 95% confidence intervals (95% CIs) for HbA1c by study design: (a) Cross-sectional studies; (b) Cohort studies.

C-peptide (+) indicates the detectable C-peptide group, while **C-peptide (-)** indicates the undetectable C-peptide group. **SD** refers to standard deviations.

The **green squares** represent individual study estimates of the SMD, with their sizes reflecting the weight of each study in the meta-analysis. **Horizontal lines** depict the 95% CIs for each study's SMD, while the **diamonds at the bottom** represent the overall pooled estimates of the SMD along with its 95% CIs.

3.2. Daily Insulin Dose

The association between C-peptide and daily insulin dose was assessed by 20 studies, including 6 cohorts^{14-18,21} and 14 cross-sectional studies^{2,4,8,18,20,22,23,26-29,31,32,34}, using different statistical approaches. Eleven^{2,8,16-18,20,22,23,27,33} out of the twenty studies found a significant crude reverse association between C-peptide and daily insulin dose; Harsunen et al.² reported that the findings remained significant after adjusting for potential confounders, including age at onset, T1DM duration, BMI, and gender [β (95% CI): - 0.09 (- 0.11 to - 0.07)]

Overall, 13 studies ^{4,8,15,17,18,20,22,23,26,29,31,32,34} provided data on the size of mean differences for daily insulin dose between C-peptide groups, of which 7 studies ^{8,17,18,20,22,23,32} consistently found a significantly lower daily insulin dose in the detectable group. Based on our pooled analysis of SMD, there was a significant difference between detectable and undetectable C-peptide groups in this regard [pooled SMD (95% CI): - 0.41 (- 0.65 to - 0.18), $I^2 = 83\%$, $p\text{-value} < 0.001$] (Table 4 and Figure 3b).

3.3. Hypoglycemia

The association between C-peptide and hypoglycemia, in terms of any occurrence of hypoglycemic events (any (n = seven studies ^{15,16,22,25,29,31,36}) or severe events (n = five studies ^{7,15,20,35,37}) or percentage of time spent in the hypoglycemic range (n = four studies ^{8,14,21,26}), was evaluated by 15 studies; one study assessed both severe and any events of hypoglycemia ¹⁵.

Regarding hypoglycemic events (any and severe cases), seven studies ^{7,15,16,20,22,25,37} found significant findings, all suggesting that the detectable group experienced hypoglycemic events less frequently; however, only two studies ^{16,20} accounted for confounding factors. Sorensen et al. ²⁰ found that individuals with T1DM in detectable C-peptide group significantly had lower odds for severe hypoglycemia after accounting for the confounding effect of T1DM duration, HbA1c level, and insulin dose [adjusted OR (95% CI): 0.4 (0.14 to 0.91)]. Jeyam et al. ¹⁶ conducted a cohort study showing that individuals with T1DM in detectable C-peptide group compared to others less frequently developed hypoglycemic events during a 5-year follow-up [Hazard ratio (HR) (95% CI): 0.35 (0.16 to 0.76)].

However, none of the four studies ^{8,14,21,26} performing glucose monitoring detected a significant difference in the percentage of time spent in the hypoglycemic range between the two study groups (Table 4).

3.4. DKA


Three cohort studies, two retrospective cohort ^{17,38} (RC) and one prospective cohort ¹⁶ (PC), with mean follow-up duration of 5.2 to 15 years, compared the incidence of DKA events between individuals with T1DM in detectable and undetectable C-peptide groups. Lee et al. ³⁸ and Jeyam et al. ¹⁶ studies found a lower risk of DKA in the detectable group; Jeyam et al. ¹⁶ considered confounding variables such as HbA1c, age at onset, sex, and T1DM duration (adjusted HR (95%

CI): 0.44 (0.29 to 0.67)). The RC study by Suh et al.¹⁷ failed to detect a significant finding in this regard in their longitudinal analysis, but their cross-sectional analysis of data from the follow-up visit revealed that detectable C-peptide is associated with lower odds of DKA [crude OR (95% CI): 0.21 (0.06 to 0.74)].


3.5. Other glycemic control indicators

Most studies reported that C-peptide levels were directly associated with the percentage of time in the normoglycemic range (seven studies^{8,14,18,21,28,32,39} out of eight studies^{8,14,18,21,26,28,32,39}), and inversely associated with serum glucose levels (five studies^{8,14,18,25,32} out of six studies^{8,14,18,25,26,32}), glucose SD (two out of two studies^{18,25}), CV (six studies^{8,18,21,25,28,39} out of seven studies^{8,18,21,25,26,28,39}), and MAGE (one out of one¹⁸) (Table 4).


Furthermore, four out of five studies^{8,14,25,26,32} (one cohort¹⁴ and three cross-sectional studies^{8,25,32}) reported a significant inverse association between C-peptide levels and the percentage of time in the hyperglycemic range. However, none of them adjusted for potential confounding factors (Table 4).




Journal Pre-proof




b



c



d



e

Figure 3: Pooled estimations of standardized mean difference (SMD) or odds ratios (OR) with 95% confidence intervals (95% CIs) for: (a) HbA1c; (b) Insulin dose; (c) Retinopathy; (d) Nephropathy; and (e) Neuropathy.

C-peptide (+) indicates the detectable C-peptide group, while **C-peptide (-)** indicates the undetectable C-peptide group. **SD** refers to standard deviations.

The **green squares** represent individual study estimates of SMD or OR, with their sizes reflecting each study's weight in the meta-analysis. **Horizontal lines** show the 95% CIs for each estimate. The **diamonds at the bottom** represent the overall pooled estimates of SMD or OR and their 95% CIs.

3.6. Microvascular Complications

A total of 16 studies investigated the association between C-peptide status/levels and microvascular complications overall ($n = 2$)^{7,13} or separately by its subtype, including retinopathy ($n = 16$)^{2,4,5,12,15-17,22,25,29,30,34,38,40-42}, nephropathy ($n = 6$)^{2,15,17,29,30,40} and its related renal conditions (CKD stage 3 ($n = 2$)^{16,34}, albuminuria [micro ($n = 5$)^{2,4,22,29,34}, macro ($n = 1$)³⁴, micro or macro ($n = 1$)³⁸], and renal failure ($n = 1$)³⁴ and/or neuropathy ($n = 6$)^{4,15,17,29,34,38}).

Four^{2,12,16,25} out of fifteen studies^{2,4,5,12,15-17,22,25,29,30,34,38,40-42} on the association between C-peptide and retinopathy found a significant finding; two of them were the largest studies included in our review, conducted by Jeyam et al.¹⁶ in 2021 (sample size = 5,732) and Harsunen et al.² in 2023 (sample size = 3,984), both of which adjusted for confounders. Based on our meta-analysis of both crude and adjusted OR, detectable C-peptide was significantly associated with lower odds of retinopathy in T1DM [pooled crude OR (95% CI): 0.53 (0.41 to 0.69), $I^2 = 65\%$, p .value < 0.001], [pooled adjusted OR (95% CI): 0.70 (0.56 to 0.87)] (Table 4 and Figures 4 and 3c).




Figure 4: Pooled analysis of adjusted odds ratios (OR) and 95% confidence intervals (95% CIs) for retinopathy. Each **blue square** represents the OR for an individual study, with the size reflecting the study's weight in the meta-analysis. **Horizontal lines** indicate the 95% CIs for each study's OR. The **diamonds at the bottom** represent the overall pooled estimates of the OR, along with their 95% CIs for both the fixed effects model and the random effects model.

A significant association between C-peptide levels and nephropathy was detected in two cross-sectional studies conducted by Gabbay et al.²⁹ [adjusted OR (95% CI): 0.4 (not reported)] and Harsunen et al.² [adjusted OR (95% CI): 0.61 (0.38 to 0.96)]. Regarding the association of C-peptide level with CKD stage 3 and renal failure, only Jeyam et al.¹⁶ observed a significant OR in their cross-sectional analysis, suggesting the presence of a direct association [adjusted OR (95% CI): 1.95 (1.25 to 3.05)]; however, they failed to confirm this significant association in their longitudinal analysis. Of six studies^{2,4,22,29,34,38} comparing the presence of albuminuria (microalbuminuria and/or macroalbuminuria) between study groups, a significantly lower odds of "microalbuminuria" in the detectable C-peptide group was reported by Gabbay et al.²⁹ [adjusted OR (95% CI): 0.39 (0.16 to 0.92)] and Harsunen et al.² [adjusted OR (95% CI): 0.71 (0.52 to 0.97)]; none of the four studies^{4,22,34,38} reporting a non-significant association considered potential confounders (Table 4).

Our meta-analysis revealed that individuals with T1DM in the detectable C-peptide group had lower odds of nephropathy and its correlated renal conditions than those in the undetectable C-peptide group [pooled crude OR (95% CI): 0.62 (0.55 to 0.70), I² = 19%, p.value < 0.001] (Figure 3d).

None of the six studies (two cohort ^{17,38} and four cross-sectional ^{4,15,29,34}) found a significant association between C-peptide levels and neuropathy complications [pooled crude OR (95%CI): 0.92 (0.65 to 1.31), I² = 0%, p.value: 0.31] (Figure 3e).

3.7. Macrovascular complications

Five studies (including four cross-sectional studies ^{2,4,15,43} and one cohort study ¹³) assessed the association between C-peptide levels and macrovascular complications. The cross-sectional study conducted by Harsunen et al. ² found a significant association, revealing that individuals with T1DM in the detectable C-peptide group had a 40% lower risk of macrovascular complications [crude OR (95% CI): 0.60 (0.44 to 0.83)]. In contrast, Irlouzadian et al. ⁴³ found that C-peptide levels were significantly associated with an increased risk of CAD among patients with T1DM [crude OR (95% CI): 1.62 (1.22 to 2.16)]. This association remained significant after adjusting for factors such as age, sex, BMI, diabetes duration, HbA1c, and fasting blood sugar [adjusted OR (95% CI): 1.65 (1.24 to 2.19)] (Table 4).

4. Discussion

This systematic review includes a total of 38 articles investigating the relationship between C-peptide status and critical parameters in individuals with T1DM. Individuals with T1DM in the detectable C-peptide group exhibited better glycemic control, as evidenced by their lower HbA1c levels and a reduced daily insulin requirement. Regarding microvascular complications, detectable C-peptide was associated with a lower risk of retinopathy and nephropathy; however, there was no association between detectable C-peptide levels and neuropathy.

Research indicates that T1DM can exhibit distinct progression patterns depending on the immunogenetic phenotypes of individuals with T1DM ⁴⁵. Young individuals with high immunogenetic risk tend to manifest the disease earlier in life and experience rapid progression toward absolute insulin deficiency ⁴⁶. However, some cases might maintain a level of insulin secretion throughout adulthood, potentially contributing positively to the course of the disease ⁴⁷.

Pancreatic islet beta cells produce proinsulin, which is cleaved into insulin and C-peptide, both of which are released into the portal circulation in an equimolar ratio ^{48,49}. Unlike insulin, which is mostly eliminated during its first pass through the liver with a half-life of 3 to 10 minutes ⁵⁰, C-peptide avoids initial metabolism and has a half-life of approximately 30 minutes, making it a

reliable indicator of beta cell function in clinical settings ⁵¹. In individuals with T1DM, C-peptide levels are significantly decreased or nearly absent ⁵². Decreases in these parameters indicate disease severity and increased reliance on exogenous insulin therapy for blood glucose management ⁵³. Additionally, a sudden increase in C-peptide levels has been associated with the remission phase of T1DM, a unique phase associated with the reduced risk of microvascular complications ⁵⁴.

Although C-peptide has long been considered a by-product of insulin secretion, a growing body of evidence suggests that it has beneficial biological properties. These potential beneficial effects have been shown in animal and human kidney and mesangium cell lines. Further investigation is warranted to explore these impacts within human clinical environments. While the current focus of T1DM therapy revolves around the administration of external insulin, there is a hypothesis that the absence of simultaneous administration of C-peptide may result in a physiological gap and potentially play a role in the emergence of long-term complications associated with the disease ¹⁰.

Although there were some inconsistencies in the included studies regarding glycaemic control, our meta-analysis showed lower HbA1C and insulin doses in individuals with T1DM in the detectable C-peptide group. In addition, most studies have indicated that individuals in the detectable C-peptide group tend to have longer periods of normoglycemia. Remarkably, these advantages have been noted alongside a decrease in hypoglycemia occurrence. This result demonstrates that undetectable C-peptide levels can lead to increased reliance on external insulin, leading to the occurrence of both hyperglycemia and hypoglycemia. Consequently, more stringent disease management strategies are required for these individuals.

Our study showed that individuals with T1DM in the detectable C-peptide group experienced fewer microvascular complications. The longer periods of normoglycemia in these individuals result in a lower rate of non-enzymatic glycosylation, resulting in less endothelial damage and fewer microvascular complications ⁵⁵. Nevertheless, over the last decade, research has revealed that this hypothesis fails to account for all the advantageous impacts of C-peptide in this context ⁵⁶. Numerous studies have presented persuasive findings indicating that C-peptide's biological functionality can safeguard against microvascular leakage by obstructing vascular endothelial growth factor (VEGF)-induced microvascular permeability. This inhibition occurs through the suppression of reactive oxygen species (ROS)-mediated intracellular processes, preservation of

VE-cadherin integrity, and prevention of stress fiber formation⁵⁶. However, most of the studies included did not adjust for diabetes duration, which may limit our understanding of C-peptide's role in glycemic control and the development of chronic complications in individuals with T1DM.

5. Conclusions

This study represents the first systematic review exploring the correlation between C-peptide levels and critical parameters in individuals with T1DM. Individuals who exhibit detectable C-peptide levels may experience enhanced control over their glycemic parameters, including reduced HbA1c levels, decreased daily insulin requirements, and decreased likelihood of DKA. Moreover, the presence of detectable C-peptide likely correlates with a lowered risk of hypoglycemic incidents. Detectable C-peptide is anticipated to reduce the risk of complications such as retinopathy, nephropathy, and related renal disorders. By addressing the physiological gap arising from insulin therapy regimens, C-peptide may open up novel avenues for enhancing the quality of life and long-term outcomes of individuals with T1DM. To comprehensively understand the direct effects of C-peptide, further investigations encompassing both basic and clinical studies are strongly encouraged.

Funding: This research received no external funding.

Acknowledgments: Researchers appreciated the Clinical Research Development Units of Kamali and Rajaee Hospitals in Alborz University of Medical Sciences.

Author Contributions: Conceptualization, M.S.1 and H.R.; methodology, M.S.1, and H.R.; software: M.S.1, P.S., and S.S.; validation, M.S.1, and H.R.; formal analysis, P.S. and M.S.1; investigation, M.S.1, T.S., R.F., and S.S.; resources, M.S.1, T.S., S.S., and R.F.; data curation, M.S.1 and M.S.2; writing original draft preparation, H.R., M.S.1, E.J., and A.T.; writing review and editing, M.S.1, S.S., and M.S.2; visualization P.S and MS1; supervision, H.R. and M.S.1; project administration, M.S.1. All authors have read and agreed to the published version of the manuscript.

Disclosure statement: The authors declare no conflicts of interest.

References:

1. Katsarou A, Gudbjörnsdottir S, Rawshani A, et al. Type 1 diabetes mellitus. *Nature reviews Disease primers*. 2017;3(1):1-17.
2. Harsunen M, Haukka J, Harjutsalo V, et al. Residual insulin secretion in individuals with type 1 diabetes in Finland: longitudinal and cross-sectional analyses. *The Lancet Diabetes & Endocrinology*. 2023;
3. Wang L, Lovejoy NF, Faustman DL. Persistence of prolonged C-peptide production in type 1 diabetes as measured with an ultrasensitive C-peptide assay. *Diabetes care*. 2012;35(3):465-470.
4. Keenan HA, Sun JK, Levine J, et al. Residual insulin production and pancreatic β -cell turnover after 50 years of diabetes: Joslin Medalist Study. *Diabetes*. 2010;59(11):2846-2853.
5. Grönberg A, Espes D, Carlsson P-O. Better HbA1c during the first years after diagnosis of type 1 diabetes is associated with residual C peptide 10 years later. *BMJ Open Diabetes Research and Care*. 2020;8(1):e000819.
6. McKeigue PM, Spiliopoulou A, McGurnaghan S, et al. Persistent C-peptide secretion in type 1 diabetes and its relationship to the genetic architecture of diabetes. *BMC medicine*. 2019;17:1-11.
7. Kuhlreiber W, Washer S, Hsu E, et al. Low levels of C-peptide have clinical significance for established Type 1 diabetes. *Diabetic Medicine*. 2015;32(10):1346-1353.
8. Rickels MR, Evans-Molina C, Bahnson HT, et al. High residual C-peptide likely contributes to glycemic control in type 1 diabetes. *The Journal of clinical investigation*. 2020;130(4):1850-1862.
9. Jones A, Hattersley A. The clinical utility of C-peptide measurement in the care of patients with diabetes. *Diabetic medicine*. 2013;30(7):803-817.
10. Ryk A, Łosiewicz A, Michalak A, Fendler W. Biological activity of c-peptide in microvascular complications of type 1 diabetes—time for translational studies or back to the basics? *International Journal of Molecular Sciences*. 2020;21(24):9723.
11. Huang Y, Wang Y, Liu C, et al. C-peptide, glycaemic control, and diabetic complications in type 2 diabetes mellitus: a real-world study. *Diabetes/Metabolism Research and Reviews*. 2022;38(4):e3514.
12. Nakanishi K, Watanabe C. Rate of β -cell destruction in type 1 diabetes influences the development of diabetic retinopathy: protective effect of residual β -cell function for more than 10 years. *The Journal of Clinical Endocrinology & Metabolism*. 2008;93(12):4759-4766.
13. Panero F, Novelli G, Zucco C, et al. Fasting plasma C-peptide and micro-and macrovascular complications in a large clinic-based cohort of type 1 diabetic patients. *Diabetes care*. 2009;32(2):301-305.
14. Carr AL, Oram RA, Marren SM, McDonald TJ, Narendran P, Andrews RC. Measurement of peak C-peptide at diagnosis informs glycemic control but not hypoglycemia in adults with type 1 diabetes. *Journal of the Endocrine Society*. 2021;5(10):bvab127.
15. Cheng J, Yin M, Tang X, et al. Residual β -cell function after 10 years of autoimmune type 1 diabetes: prevalence, possible determinants, and implications for metabolism. *Annals of Translational Medicine*. 2021;9(8)
16. Jeyam A, Colhoun H, McGurnaghan S, et al. Clinical impact of residual C-peptide secretion in type 1 diabetes on glycemia and microvascular complications. *Diabetes Care*. 2021;44(2):390-398.
17. Suh J, Lee HI, Lee M, et al. Insulin Requirement and Complications Associated With Serum C-Peptide Decline in Patients With Type 1 Diabetes Mellitus During 15 Years After Diagnosis. *Frontiers in Endocrinology*. 2022;13:869204.
18. Zhang L, Xu Y, Jiang X, et al. Impact of flash glucose monitoring on glycemic control varies with the age and residual β -cell function of patients with type 1 diabetes mellitus. *Journal of Diabetes Investigation*. 2022;13(3):552-559.
19. Grönberg A, Espes D, Carlsson P-O, Ludvigsson J. Higher risk of severe hypoglycemia in children and adolescents with a rapid loss of C-peptide during the first 6 years after type 1 diabetes diagnosis. *BMJ Open Diabetes Research and Care*. 2022;10(6):e002991.
20. Sørensen JS, Johannesen J, Pociot F, et al. Residual β -cell function 3–6 years after onset of type 1 diabetes reduces risk of severe hypoglycemia in children and adolescents. *Diabetes Care*. 2013;36(11):3454-3459.
21. Buckingham B, Cheng P, Beck RW, et al. CGM-measured glucose values have a strong correlation with C-peptide, HbA 1c and IDAAC, but do poorly in predicting C-peptide levels in the two years following onset of diabetes. *Diabetologia*. 2015;58:1167-1174.
22. Marren S, Hammersley S, McDonald T, et al. Persistent C-peptide is associated with reduced hypoglycaemia but not HbA1c in adults with longstanding Type 1 diabetes: evidence for lack of intensive treatment in UK clinical practice? *Diabetic Medicine*. 2019;36(9):1092-1099.
23. Miao H, Zhang J, Gu B, et al. Prognosis for residual islet β -cell secretion function in young patients with newly diagnosed type 1 diabetes. *Journal of Diabetes*. 2019;11(10):818-825.

24. Sugihara S, Kikuchi T, Urakami T, et al. Residual endogenous insulin secretion in Japanese children with type 1A diabetes. *Clinical Pediatric Endocrinology*. 2021;30(1):27-33.
25. Gibb FW, McKnight JA, Clarke C, Strachan MW. Preserved C-peptide secretion is associated with fewer low-glucose events and lower glucose variability on flash glucose monitoring in adults with type 1 diabetes. *Diabetologia*. 2020;63:906-914.
26. McDonald TJ, Oram RA, Shaw JA, West DJ. Postexercise Glycemic Control in Type 1 Diabetes Is Associated With Residual β -Cell Function. *Diabetes Care*. 2020;43
27. Ito A, Horie I, Miwa M, et al. Impact of glucagon response on early postprandial glucose excursions irrespective of residual β -cell function in type 1 diabetes: A cross-sectional study using a mixed meal tolerance test. *Journal of Diabetes Investigation*. 2021;12(8):1367-1376.
28. Taylor GS, Shaw AC, Smith K, et al. Capturing the real-world benefit of residual β -cell function during clinically important time-periods in established Type 1 diabetes. *Diabetic Medicine*. 2022;39(5):e14814.
29. Gabbay MA, Crispim F, Dib SA. Residual β -cell function in Brazilian Type 1 diabetes after 3 years of diagnosis: prevalence and association with low presence of nephropathy. *Diabetology & Metabolic Syndrome*. 2023;15(1):51.
30. Majaliwa ES, Muze KC, Ndayongeje J, Mfinanga SG, Mmbaga BT, Ramaiya K. Correlation of c-peptide with complications observed in children and adolescents with type 1 diabetes in Tanzania: A cross-sectional survey. *Global Pediatric Health*. 2023;10:2333794X231159790.
31. Lanctôt SO, Lovblom LE, Lewis EJ, et al. Fasted C-Peptide Distribution and Associated Clinical Factors in Adults With Longstanding Type 1 Diabetes: Analysis of the Canadian Study of Longevity in Type 1 Diabetes. *Canadian Journal of Diabetes*. 2024;48(2):89-96.
32. Liu W, Fang Y, Cai X, et al. Preserved C-peptide is common and associated with higher time in range in Chinese type 1 diabetes. *Frontiers in Endocrinology*. 2024;15:1335913.
33. Beato-Víbora PI, Tormo-García MÁ. Glycemic control and insulin requirements in type 1 diabetic patients depending on the clinical characteristics at diabetes onset. *Endocrine Research*. 2014;39(2):86-90.
34. Williams KV, Becker DJ, Orchard TJ, Costacou T. Persistent C-peptide levels and microvascular complications in childhood onset type 1 diabetes of long duration. *Journal of Diabetes and its Complications*. 2019;33(9):657-661.
35. Kristensen PL, Pedersen-Bjergaard U, Schalkwijk C, Olsen NV, Thorsteinsson B. Erythropoietin and vascular endothelial growth factor as risk markers for severe hypoglycaemia in type 1 diabetes. *European journal of endocrinology*. 2010;163(3):391-398.
36. Henriksen MM, Færch L, Thorsteinsson B, Pedersen-Bjergaard U. Long-term prediction of severe hypoglycemia in type 1 diabetes: is it really possible? *Journal of diabetes science and technology*. 2016;10(6):1230-1235.
37. Wellens MJ, Vollenbrock CE, Dekker P, et al. Residual C-peptide secretion and hypoglycemia awareness in people with type 1 diabetes. *BMJ Open Diabetes Research and Care*. 2021;9(1):e002288.
38. Lee TH, Kwon AR, Kim YJ, Chae HW, Kim HS, Kim DH. The clinical measures associated with C-peptide decline in patients with type 1 diabetes over 15 years. *Journal of Korean medical science*. 2013;28(9):1340-1344.
39. Babaya N, Noso S, Hiromine Y, et al. Relationship of continuous glucose monitoring-related metrics with HbA1c and residual β -cell function in Japanese patients with type 1 diabetes. *Scientific Reports*. 2021;11(1):4006.
40. Bhagadurshah RR, Eagappan S, Santharam RK, Subbiah S. The Impact of Body Mass Index, Residual Beta Cell Function and Estimated Glucose Disposal Rate on the Development of Double Diabetes and Microvascular Complications in Patients With Type 1 Diabetes Mellitus. *Cureus*. 2023;15(11)
41. Jensen RA, Agardh E, Lernmark Å, et al. HLA genes, islet autoantibodies and residual C-peptide at the clinical onset of type 1 diabetes mellitus and the risk of retinopathy 15 years later. *PloS one*. 2011;6(3):e17569.
42. Rajalakshmi R, Amutha A, Ranjani H, et al. Prevalence and risk factors for diabetic retinopathy in Asian Indians with young onset type 1 and type 2 diabetes. *Journal of Diabetes and its Complications*. 2014;28(3):291-297.
43. Irilouzadian R, Afaghi S, Esmaeili Tarki F, Rahimi F, Malekpour Alamadari N. Urinary c - peptide creatinine ratio (UCPCR) as a predictor of coronary artery disease in type 1 diabetes mellitus. *Endocrinology, Diabetes & Metabolism*. 2023;6(3):e413.
44. Hwang JW, Kim MS, Lee D-Y. Factors associated with c-peptide levels after diagnosis in children with type 1 diabetes mellitus. *Chonnam medical journal*. 2017;53(3):216-222.
45. team Ts, Leete P, Oram RA, et al. Studies of insulin and proinsulin in pancreas and serum support the existence of aetiopathological endotypes of type 1 diabetes associated with age at diagnosis. *Diabetologia*. 2020/06// 2020;63(6):1258-1267. doi:10.1007/s00125-020-05115-6

46. Inshaw JRJ, Cutler AJ, Crouch DJM, Wicker LS, Todd JA. Genetic Variants Predisposing Most Strongly to Type 1 Diabetes Diagnosed Under Age 7 Years Lie Near Candidate Genes That Function in the Immune System and in Pancreatic β -Cells. *Diabetes Care*. 2020/01/01/ 2020;43(1):169-177. doi:10.2337/dc19-0803
47. Oram RA, Jones AG, Besser REJ, et al. The majority of patients with long-duration type 1 diabetes are insulin microsecretors and have functioning beta cells. *Diabetologia*. 2014/01// 2014;57(1):187-191. doi:10.1007/s00125-013-3067-x
48. Strachan MWJ, Frier BM. History, Normal Physiology, and Production of Insulin. *Insulin Therapy*. Springer London; 2013:1-8.
49. Vasiljević J, Torkko JM, Knoch K-P, Solimena M. The making of insulin in health and disease. *Diabetologia*. 2020/10// 2020;63(10):1981-1989. doi:10.1007/s00125-020-05192-7
50. Kelkar S, Muley S, Ambardekar P. Insulin Pharmacodynamics, Pharmacokinetics, and Insulin Regimens. *Towards Optimal Management of Diabetes in Surgery*. Springer Singapore; 2019:275-289.
51. Landreh M, Jörnvall H. C-peptide evolution: Generation from few structural restrictions of bioactivities not necessarily functional. *FEBS Letters*. 2015/02/13/ 2015;589(4):415-418. doi:10.1016/j.febslet.2015.01.006
52. Maheshwari A. Insights on C-peptide in diabetes. *IJNMHS*. 2023/07/28/ 2023;6(2):63-65. doi:10.18231/j.ijnmhs.2023.009
53. Leighton E, Sainsbury CAR, Jones GC. A Practical Review of C-Peptide Testing in Diabetes. *Diabetes Ther*. 2017/06/01/ 2017;8(3):475-487. doi:10.1007/s13300-017-0265-4
54. Neylon OM, White M, O'Connell MA, Cameron FJ. Insulin-dose-adjusted HbA_{1c}-defined partial remission phase in a paediatric population—when is the honeymoon over? *Diabetic Medicine*. 2013/05// 2013;30(5):627-628. doi:10.1111/dme.12097
55. Wang R, Yu X, Gkousioudi A, Zhang Y. Effect of Glycation on Interlamellar Bonding of Arterial Elastin. *Exp Mech*. 2021/01// 2021;61(1):81-94. doi:10.1007/s11340-020-00644-y
56. Souto SB, Campos JR, Fangueiro JF, et al. Multiple Cell Signalling Pathways of Human Proinsulin C-Peptide in Vasculopathy Protection. *International Journal of Molecular Sciences*. 2020/01/18/ 2020;21(2):645. doi:10.3390/ijms21020645

Journal Pre-proof

Table 1: Assessment of the included studies (cohort studies)

Author	Selection			Comparability	Outcome			Total Of 9 scores
	Representativeness of exposed cohort	Selection of non-exposed cohort	Ascertainment of exposure		Assessment of outcome	Enough follow-up to occur outcomes	Adequacy of follow up	
1. Nakanishi et al.	*	*	*	*	**	*	*	9
2. Panero et al.	*	*	*	*	**	*	*	9
3. Kristensen et al.	*	*	*	*	*	-	*	7
4. Jensen et al.	*	*	*	*	**	*	*	9
5. Lee et al.	*	*	*	*	*	*	*	8
6. Beato-Vibora et al.	*	*	*	*	*	*	*	8
7. Henriksen et al.	*	*	*	*	**	*	*	9
8. Carr et al.	*	*	*	*	**	*	*	9
9. Jeyam et al.	*	*	*	*	**	*	*	9
10. Suh et al.	*	*	*	*	*	*	*	8
11. Zhang et al.	*	*	*	*	**	*	*	9
12. Grönberg et al ^b .	*	*	*	*	*	*	*	9

Table 2: Assessment of the included studies (cross-sectional studies)

Author	Selection			Comparability	Outcome		Total score Of 7 scores
	Representativeness of the sample:	Non-respondents:	Ascertainment of the exposure (risk factor):		Assessment of the outcome:	Statistical test:	
1. Keenan et al.	*	*	*	*	*	*	6
2. Sorensen et al.	*	*	*	**	*	*	7
3. Rajalakshmi et al.	*	*	*	**	*	*	7
4. Kuhlreiber et al.	*	*	*	*	*	*	6
5. Buckingham et al.	*	*	*	*	*	*	6
6. Hwang et al.	*	*	*	*	*	*	6
7. Marren et al.	*	*	*	*	*	*	6
8. Williams et al.	*	*	*	*	*	*	6
9. Miao et al.	*	*	*	**	*	*	7
10. Rickels et al.	*	*	*	*	*	*	6
11. Sugihara et al.	*	*	*	*	*	*	6
12. Gibb et al.	*	*	*	**	*	*	7
13. Gronberg et al ^a .	*	*	*	**	*	*	7
14. Taylor et al ^b .	*	*	*	**	*	*	7
15. Ito et al.	*	*	*	*	*	*	6
16. Cheng et al.	*	*	*	*	*	*	6
17. Wellens et al.	*	*	*	**	-	*	6
18. Babaya et al.	*	*	*	*	*	*	6
19. Taylor et al ^a .	*	*	*	*	*	*	6

Journal Pre-proof

20. Gabbay et al.	*	*	*	*	*	*	6
21. Harsunen et al.	*	*	*	**	*	*	7
22. Majaliwa et al.	*	*	*	*	*	*	6
23. Irilouzadian et al.	*	*	*	*	*	*	6
24. Bhagadurshah et al.	*	*	*	*	*	*	6
25. Lanctôt et al.	*	*	*	**	*	*	7
26. Liu et al.	*	*	*	*	*	*	6

Journal Pre-proof

Table 3: Characteristic of the included studies

Author	Country	Study Type	Inclusion criteria	Sample size			% Male (N)			Age at onset			Disease duration			Body mass index (Kg/m ²)		
				All	C-peptide		All	C-peptide		All	C-peptide		All	C-peptide		All	C-peptide	
					N	P		Ne	Pos.		Ne	Pos.		Neg	Pos.		Ne	Pos.
1.Nakanishi et al.	Japan	RC	Outpatients/	254	104	42	57%	63%	52%	34.0	36(1)	30(12)*	5-20	20	20	NR	NR	NR
2.Panero et al.	Italy	RC	Age <60	471	201	270	NR	55%	57%	NR	NR	NR	15.8(10.)	13.9(9.1)	7.8(9.)	NR	24.1	24.5
3.Kristensen et al.	Denmark	PC	Disease D. ≥	219	NR	NR	59%	NR	NR	25(1)	NR	NR	21(12)	NR	NR	25(3.6)	NR	NR
4.Jensen et al.	Sweden	PC	Dig. Age <	246	NR	NR	55%	NR	NR	24.9	NR	NR	NR	NR	NR	22.1(3.7)	NR	NR
5.Lee et al. 2013	S.Korea	RC	Disease D.	95	76	19	41%	NR	NR	7.8(4.)	NR	NR	NR	NR	NR	0.23(1.05)	NR	NR
6.Beato-Vibora	Spain	RC	Newly diagnosed	301	NR	NR	62%	NR	NR	27(1)	NR	NR	6.8(4.5)	NR	NR	NR	NR	NR
7.Henrikssen et al.	Denmark	PC	Age >18	98	42	91	57%	NR	NR	NR	NR	NR	32(11)	NR	NR	25(4)	NR	NR
8.Carr et al. 2021	UK	PC	EXTOD	26	NR	NR	57%	NR	NR	27.2	NR	NR	New case	NR	NR	23.5(3.1)	NR	NR
9.Jeyam et al.	UK	PC	SDRN T1BIO	5732	3571	2161	56%	54.7%	58.3%	20.8	17.0	25.9(14.4)	20.9(14.)	25.4(17.)	11.7	26.3(5.8)	26.4	26.2
10.Suh et al.	S.Korea	RC	Disease D. ≥	234	77	157	43%	NR	NR	8.3	NR	NR	NR	NR	NR	0.25(1.15)	NR	NR
11.Zhang et al.	China	RC/C	Disease D. ≥	171	112	59	43%	43%	42%	NR	NR	NR	1.6(1.9)	2.0(2)	0.7(0.)	18.6(3.4)	18.7	18.5
12.Gronberg et al.	Sweden	PC/C	Newly diagnosed	50	30	20	44%	50%	35%	10.6	11.1	10.0(2.3)	NR	NR	NR	-0.07(1.26)	0.08	-0.29
13.Keenan et al.	US	CS	Disease D. ≥	382	126	256	48.4%	42.6%	51%	11(6.)	10.9	11.1(6.2)*	56.2(5.8)	56.4(5.7)	56.0	26.0(5.1)	26.7	25.9
14.Sorensen et al.	Denmark	CS	Dig. Age <	342	250	92	51%	51%	49%	9.2(3.)	8.4(3.)	9.0(2.6)*	4.2(0.9)	4.4(0.9)	3.9(0.)	NA	NA	NA
15.Rajalakshmi	India	CS	Dig. Age <	150	NR	NR	56%	NR	NR	16.8	NR	NR	12.4(7.4)	NR	NR	22.1(3.6)	NR	NR
16.Kuhtr eiber et al. 2015	US /Ca	CS	-	1272	NR	NR	43%	NR	NR	20.6(4.)	NR	NR	19(14.8)	NR	NR	NR	NR	NR
17.Buckingham	US	CS	Newly diagnosed	67	NR	NR	NR	NR	NR	NR	NR	NR	New case	NR	NR	NR	NR	NR
18.Hwang et al. 2017	Korea	CS	Newly diagnosed	34	27	7	44%	11%	33%	10(4.)	9(4.)	13.6(3.6)*	3(NR)	3(NR)	3(NR)	16.1(2.8)	15.5	18.7
19.Marr en et al. 2019	UK	CS	Dig. Age <	221	151	70	52%	57%	41%*	8.9(8.)	6.1(7.)	15.1(7.2)*	13.0(11.)	13.3(11.)	12.6(10.)	23.9(4.2)	23.3	25.2
20.Williams et al.	US	CS	Dig. Age <	185	167	18	49%	49%	50%	<17	8.1(4.)	9.7(4.2)	42.9(6.7)	43.0(6.7)	42.1	28.2(5.1)	28.4	26.7
21.Miao et al. 2019	China	CS	C-peptide	110	68	42	43%	36%	55%*	20.2(5.)	19.2(5.)	22.8(5.9)	2.5(1.8)	2.3(1.7)	2.7(1.)	19.4(4.4)	19.1	19.8
22.Rickels et al.	US	CS	Age 18 to	63	15	48	46%	53%	52%	NR	NR	NR	7.4(6.3)	13(9)	5.6(3.)	24.2(2.9)	25(3)	24(3)
23.Sugihara et al.	Japan	CS	Dig. Age <	576	NR	NR	40%	NR	NR	5	NR	NR	1.1	NR	NR	NR	NR	NR
24.Gibb et al.	UK	CS	Disease D. ≥	290	201	89	56%	54%	60%	17.4	15(1)	23(10.3)	21.9(13.)	25(14.)	15(11)	26.7(4.6)	26.6	27.2
25.Gronberg et al.	Sweden	CS	Age < 25	73	45	28	55%	64%	39%*	5.1(3.)	5.0(2.)	5.2(3.1)	12.3(2.4)	12.7(2.6)	11.8	0.52(1.06)	0.28	0.90

26.Taylor et al ^b .	UK	CS	Disease D. ≥	30	11	19	53 %	45 %	58 %	18.2	13.27	21 (9.2)*	19.9 (13.	26.8 (13.	16.0	25.2 (3.7)	25.6	24.9
27.Ito et al. 2020	Japan	CS	Age >16	34	17	17	47 %	47 %	59 %	NR	NR	NR	8.1 (9.1	11.1 (9.2	5.1 (8.	22.5 (4.5)	22.5	22.6
28.Chen g et al.	China	CS	Disease D.	109	67	42	40 %	45 %	33 %	19.0	17.0	27.5(25.5)	13 (3.7	13.0 (3.7	12.0	20.6 (2.5)	20.6	20.7
29.Well ens et al.	Netherlands	CS	Disease D. ≥	509	NR	NR	41 %	NR	NR	12 (8.	NR	NR	19 (13.	NR	NR	25.0 (3.59)	NR	NR
30.Babaya et al.	Japan	CS	Non pregna	19	NR	NR	32 %	NR	NR	NR	NR	NR	8.8 (9.6	NR	NR	21.7 (3.8)	NR	NR
31.Taylor et al ^a .	UK	CS	Disease D ≥	66	34	32	56 %	53 %	53 %	20 (1	18 (1	22 (12.5)	21.5 (12.	25 (11)	17.9	25.4 (3.3)	25.4	25.5
32.Gabb ay et al.	Brazil	CS	Disease D. ≥	138	56	82	59 %	57 %	60 %	9.0 (0.	8.7 (5.	10.2 (5.6)*	12.0 (0.5	14.7 (6.7	11.4	23.0 (0.3)	23.1	23.5
33.Hars unen et	Finland	CS	FinnDiane	3984	3208	776	51 %	50 %	55 %	13.5	12.6	18.6 (9.8)*	21.6 (13.	23.3 (12.	12.1	24.7 (3.7)	24.8	24.4
34.Majal iwa et	Tanzania	CS	Age: 0 to 20	281	200	81	49 %	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR
35.Irilou zadian et	Iran	CS	Disease D. ≥	279	NR	NR	50 %	NR	NR	NR	NR	NR	43.4 (4.1	NR	NR	26.5 (4.7)	NR	NR
36.Bhag adurshah	India	CS	Disease D. ≥	113	NR	NR	55 %	NR	NR	15 (3.	NR	NR	16 (6.7	NR	NR	22.04 (3.9)	NR	NR
37.Lanct ôt et al.	Canada	CS	Disease D. >	74	26	48	55 %	58 %	54 %	NR	10 (6.	10 (8.1)	54 (4.4	54 (3.7	54 (4.	NR	26.5	26.7
38.Liu et al. 2024	China	CS	fCP >1,500	534	238	296	46 %	42 %	49 %	36 (2	30 (2	38 (20.0)	9.0 (13.	15.0 (8.	4.0	22.8 (3.3)	23.1	22.

ACEIs: Angiotensin Converting Enzyme Inhibitors; ARBs: Angiotensin Receptor Blockers; BS: Blood sugar; Ca: Canada; CS: Cross-sectional; DISS: Diabetes Incidence Study in Sweden; Dig. Age: Diagnosis age; Disease D.: Disease Duration; S.Korea: South Korea; EDC: Pittsburgh Epidemiology of Diabetes Complications; FinnDiane: The Finnish Diabetic Nephropathy; EXTOD: Exercise in Type 1 Diabetes; JSGIT: The Japanese Study Group of Insulin Therapy for Childhood and Adolescent Diabetes; Neg.: Negative (Undetectable group); NR: Not reported; PC: prospective cohort; Pos.: Positive (Detectable group); RC: Retrospective cohort; SDRNT1BIO: The Scottish Diabetes Research Network Type 1 Bioresource; * P Value <0.05, † SDS, ‡ Panel study

Journal Pre-proof

Table 4: Association of detectable C-peptide levels with glycemic control and chronic complications in patients with Type 1 Diabetes mellitus

I d	Author	Type study	C-peptide method	C-peptide test	FU D.	IV Cut-P. Pmol/L	Outcomes	Sign.	Direction	Effect size for C- peptide (95% CI) [Detectable vs. Undetectab le]	Confounder s
1	Nakanishi et al.	RC	RIA	fCP/PCP	5-20	fCP > 17 or PCP > 33	HbA1c at 5 & 10 Yrs. Follow up HbA1c at 15 Yrs. Follow up HbA1c at 20 Yrs. Follow up Retinopathy at Yrs. Follow up Retinopathy at 10 Yrs. Follow up Retinopathy at 15 Yrs. Follow up Retinopathy at 20 Yrs. Follow up	NS + + NS + + + +	NA ↓ ↓ NA ↓ ↓ ↓ ↓	- MD= - 0.44 (-0.76 to -0.12) MD= - 0.59 (-0.93 to -0.25) - HR= 0.70 (0.53 to 0.93) HR= 0.65 (0.48 to 0.88) HR= 0.61 (0.37 to 0.96)	None // // Sex, HTN, HbA1c, time to insulin therapy, age at onset // // //
2	Panero et al.	RC	NR	fCP	4.5	60	HbA1c at baseline Microvascular complications Macrovascular complications	NS + NS	NA ↓ NA	- OR= 0.59 (0.37 to 0.94) - -	None Age, sex, Duration of diabetes, HbA1c, CVD, and HTN //
3	Kristensen et al.	PC	RIA	rCP	1	10	Severe hypoglycemia	NS	NA	-	None
4	Jensen et al.	PC	NR	NR	15	130	Retinopathy	NS	NA	-	HbA1c, HTN, HLA and islet autoantibod ies
5	Lee et al.	RC	RIA	fCP	15	50	Retinopathy Peripheral Neuropathy Albuminuria	NS NS NS +	NA NA NA ↓	- - - Risk Diff. = 26.3 %	None // // //

Journal Pre-proof

							History of DKA			(1.5% to 43%)	
6	Beato-Vibora et al	RC	CLIA	fCP /GST	6(4.8)	NR	Insulin dose	+	↓	P value <0.05	None
7	Henriksen et al.	PC	NR	NR	12	10	Hypoglycemia	NS	NA	-	None
8	Carr et al.	PC	NR	MMTT	1	100	HbA1c	NS	NA	-	None
							Inulin dose	NS	NA	-	//
							% Time in normoglycemia	+	↑	$\beta = 2.39\%$ (0.51-4.26)	//
							glucose SD	+	↓	$\beta = -0.14$ (-0.25 to -0.023)	//
							% Time in hyperglycemia	NS	NA	$\beta = -2.64$ (-4.87 to -0.41)	//
							% Time in hypoglycemia			-	
9	Jeyam et al.	PC	ECLIA	rCP	5.2	> 200 vs. < 5	HbA1c at Baseline	+	↓	$\beta = -6.41$, (-7.9 to -4.9)	None
							HbA1c at Fu.	+	↓	$\beta = -4.90$ (-6.2 to -3.6)	//
							Insulin dose	+	↓		
							DKA during Fu.	+	↓	$\beta = -0.04$ (-0.0 to -0.03)	HbA1c, Age at onset, Sex, and Duration of diabetes
							Hypoglycemia during Fu.	+	↓	HR= 0.44 (0.29 to 0.67)	//
							Retinopathy at Baseline	+	↑		
							Retinopathy at Fu.	NS	NA	HR= 0.35 (0.16 to 0.76)	//
							CKD Stage 3 at Baseline			OR= 0.66 (0.51 to 0.86)	//
							CKD at Fu.			OR= 0.51 (0.35 to 0.74)	//
										OR= 1.95 (1.25 to 3.05)	
										-	
10	Suh et al.	RC	RIA	PCP	15	500	HbA1c	NS	NA	-	None
							Insulin dose	+	↓	MD= -0.14 (-0.22 to -0.07)	//
							Insulin dose [CS Analysis]	+	↓		//
							DKA	NS	NA		//
								+	↓		//

Journal Pre-proof

							DKA [CS Analysis]	NS	NA	MD= -0.14 (-0.23 to -0.05)	//
							Retinopathy	NS	NA		//
							Neuropathy	NS	NA	-	//
							Nephropathy			OR= 0.21 (0.06 to 0.74)	
										-	
										-	
										-	
1 1	Zhang et al.	RC	CLIA	MMTT	0.5	200	HbA1c at baseline	NS	NA	-	None
							HbA1c changes in 6 months	+	↓	P value <0.05	//
							% Time normoglycemia	+	↑	R= + 0.359	//
							mean glucose	+	↓	R= - 0.218	//
							glucose SD	+	↓	R= - 0.418	//
							MAGE	+	↓	R= - 0.393	//
							Glucose CV	+	↓	R= - 0.435	//
							Insulin dose at baseline	+	↓	MD= -0.15 (-0.23 to -0.07)	//
1 2	Gronberg et al ^b .	PC	FIA	MMTT	6	30	HbA1c at diagnosis and FU. visits	NS	NA	-	None
							Insulin dose at diagnosis and FU. visits	NS	NA	-	//
1 3	Keenan et al.	CS	RIA	MMTT	NA	30	HbA1c	+	↓	MD: -0.40 (-0.61 to -0.19)	None
							Insulin dose	NS	NA		//
							Retinopathy	NS	NA	-	//
							Microalbuminuria	NS	NA	-	//
							Neuropathy	NS	NA	-	//
							CVD	NS	NA	-	//
1 4	Sorensen et al.	CS	FIA	MMTT	NA	40	HbA1c [C-peptide >200 vs. < 40]	+	↓	Diff in % = -12.9% (-17.9 to -7.5)	Sex, age, pubertal status, Duration of diabetes, insulin admin technique
							HbA1c [C-peptide >40-200 vs. < 40]	+	↓	Diff in % = -5.3% (-9.1 to -1.3)	

Journal Pre-proof

							Insulin dose [C-peptide :>200 vs. < 40]			Diff in % = -21.1% (-30.5 to -10.4)	//
							Severe hypoglycemia			OR= 0.4 (0.14 to 0.91)	// & HbA1c level, and insulin dose
15	Rajalakshmi et al.	CS	EL	fCP /PCP	NA	NR	Retinopathy	NS	NA	-	None
16	Kuhtreiber et al.	CS	ELISA	fCP	NA	10	HbA1c	+	↓	P value < 0.05	None
							Severe hypoglycemia	+	↓	//	//
							Microvascular complications	+	↓	//	//
17	Buckingham et al.	CS [‡]	NR	MMTT	2	NA	HbA1c	+	↓	P value < 0.05	None
							% Time in normoglycemia	+	↑	β= 0.53 (0.23 to 0.83)	//
							Glucose CV	NS	NA	β= -1.19 (-1.69 to -0.69)	//
							% Time in hypoglycemia			-	//
18	Hwang et al.	CS	IA	fCP	NA	198	HbA1c	NS	NA	-	None
19	Marren et al.	CS	CLIA	MMTT	NA	20	HbA1c	NS	NA	-	None
							Hypoglycemia	+	↓	HR = 0.79 (NR)	//
							Insulin dose	+	↓	MD= -0.13 (-0.20 to -0.06)	//
							Retinopathy	NS	NA		//
							Microalbuminuria	NS	NA	-	//
										-	
20	Williams et al.	CS	ELISA	fCP	NA	1.15	Insulin dose	NS	NA	-	None
							Retinopathy	NS	NA	-	//
							Microalbuminuria	NS	NA	-	//
							Macroalbuminuria	NS	NA	-	//
							CKD stage 3	NS	NA	-	//
							Renal failureDS	NS	NA	-	//
							Polyneuropathy				

Journal Pre-proof

2 1	Miao et al.	CS	IA	OGTT	NA	600	HbA1c	+	↓	MD= -1.39 (-2.07 to - 0.71)	None
							Insulin dose	+	↓		//
							Mean HbA1c during proceeding years (median 32.5 months)	+	↓	MD= -0.19 (-0.27 to - 0.11) OR= 0.41 (0.24 to 0.68)	HLA, age at onset, sex, family history of T1DM, DKA, BMI
2 2	Rickels et al.	CS	NR	MMTT	NA	0.007	HbA1c	NS	NA	-	None
							Insulin dose	+	↓	MD= -0.20 (-0.32 to - 0.08)	//
							Mean glucose	+	↓		//
							% Time in normoglycemia	+	↑	R = -0.356	//
								+	↓	r = +0.456	//
							% Time in hyperglycemia	+	↓	r = -0.376	//
								NS	NA	r = -0.258	//
							Glucose CV			-	
							% Time in hypoglycemia				
2 3	Sugihara et al.	CS	ECLIA	rCP	NA	NR	HbA1c	+	↓	R = -0.16	None
2 4	Gibb et al.	CS	IA	rCP	NA	10	HbA1c	NS	NA	-	None
							Retinopathy	+	↓	OR= 0.42 (0.25 to 0.70)	//
							Glucose SD > 4.0 mmol/l	+	↓		//
								+	↓	OR= 0.32 (0.16to0.63)	//
							Glucose CV > 40.3% [med]	+	↓		//
							Interquartile range glucose > 5.6 mmol/l [med]	+	↓	OR= 0.30 (0.17 to 0.52)	//
								+	↓		//
							Hypoglycemia > 9 times / 2 Weeks [med]			OR= 0.40 (0.21- 0.75)	
							% Time in hyperglycemia > 4% [med]			OR= 0.29 (0.15- 0.54)	
										OR= 0.32 (0.17- 0.60)	
2 5	Gronberg et al ^a .	CS	ELISA	rCP	NA	1.17	HbA1c	NS	NA	-	None
							Retinopathy	NS	NA	-	//
2 6	Taylor et al ^b .	CS	ECLIA	MMTT	NA	3	HbA1c	NS	NA	-	None
							Insulin dose	NS	NA	-	//
							Mean glucose	NS	NA	-	//

Journal Pre-proof

							% Time in normoglycemia	NS	NA	-	//
								NS	NA	-	//
							% Time in hyperglycemia	NS	NA	-	//
								NS	NA	-	//
							Glucose CV				
							% Time in hypoglycemia				
27	Ito et al.	CS	ECLusys	MMTT	NA	100	HbA1c	NS	NA	-	None
							Insulin dose	+	↓	MD= -17.3 (-28.8 to -5.8)	//
28	Cheng et al.	CS	CLIA	fCP/PCP	NA	16.7	HbA1c	+	↓	MD= -7.80 (-15.29 to -0.31)	None
							Hypoglycemia	+	↓		//
							Severe hypoglycemia	NS	NA	OR= 0.29 (0.12 to 0.67)	//
								NS	NA		//
							Insulin dose	NS	NA	-	//
							Retinopathy	NS	NA	-	//
								NS	NA	-	//
							Proliferative retinopathy	NS	NA	-	//
								NS	NA	-	//
							Nephropathy	NS	NA	-	//
								NS	NA	-	//
							Sever nephropathy	NS	NA	-	//
								NS	NA	-	//
							peripheral neuropathy			-	
							macrovascular complications			-	
29	Wellens et al.	CS	IRMA	fCP	NA	3.8	Impaired awareness of hypoglycemia	+	↓	OR= 0.51 (0.26 to 0.98)	Age at onset, BMI, and T1DM complications
							Severe hypoglycemia (self-reported)	+	↓	OR= 0.64 (0.38 to 1.08)	None
30	Babaya et al.	CS	NR	fCP	NA	NR	Glucose CV	+	↓	R= - 0.64	None
31	Taylor et al.	CS	NR	UCPCR	NA	> 200 vs. < 1	HbA1c	NS	NA	-	None
							Insulin dose	NS	NA	-	//
							% Time in Nocturnal normoglycemia	+	↑	MD= 18.00 (6.50 to 29.50)	//
								+	↑		//
							% Time in PP normoglycemia	+	↓	MD= 17.00 (4.35 to 29.65)	//

Journal Pre-proof

							Nocturnal CV (mmol/L)			MD= -0.40 (-0.63 to -0.16)	
3 2	Gabbay et al.	CS	ELISA	fCP	NA	1.15	HbA1c	NS	NA	-	None
							Insulin dose	NS	NA	-	//
							Hypoglycemia	NS	NA	-	//
							Nephropathy	+	↓	OR= 0.4 (NR)	Diabetes duration and HbA1c.
							Microalbuminuria	+	↓	OR= 0.39 (0.16 to 0.92)	//
							Neuropathy	NS	NA		None
							Retinopathy	NS	NA	-	//
										-	
3 3	Harsunen et al.	CS	UIA	rCP	NA	≥ 20	HbA1c	+	↓	β= - 4.97 (-6.28 to -3.67)	Age at onset, sex, duration of diabetes, and BMI
							Insulin dose	+	↓		
							Nephropathy	+	↓	β= - 0.09(-0.11 to -0.07)	//
							microalbuminuria	+	↓		
							Retinopathy	+	↓	OR= 0.61 (0.38 to 0.96)	// & HbA1c and its variability, hypertension, smoking, and eGFR
							Hard CVD events	+	↓	OR= 0.71 (0.52 to 0.97)	
										OR= 0.55 (0.34 to 0.89)	//
										OR= 0.60 (0.44 to 0.83)	None
3 4	Majaliwa et al.	CS	ELISA	fCP	NA	200	HbA1c	NS	NA	-	None
							Nephropathy	NS	NA	-	//
							Retinopathy	NS	NA	-	//
3 5	Irilouzadian et al.	CS	CIA	UCPCR	NA	NR	Coronary artery disease	+	↓	OR= 1.54 (1.16 to 2.05)	Adjusted for age, sex, BMI, diabetes duration, HbA1c, FBS, active smoking, alcohol consumption, HTN, Lipid profile, eGFR, uric acid, albumin, creatinine

Journal Pre-proof

											and albuminuria .
3 6	Bhagadurs hah et al.	CS	ECLIA	rCP	NA	NR	Retinopathy Nephropathy	NS NS	NA NA	- -	None //
3 7	Lanctôt et al.	CS	IA	fCP	NA	33	HbA1cInsulin dose Hypoglycemic events in past month (per week)	NS NS NS	NA NAN A	- - -	None // //
3 8	Liu et al.	CS	ECLIA	fCP/MM TT	NA	10	Diabetic retinopathy Insulin dose HbA1c > 7% eGFR % Time in normoglycemi a Mean glucose % Time in hyperglycemia	+ + NS NS + + + +	↓ ↓ NA NA ↑ ↓ ↓	OR= 0.63 (0.41 to 0.97) OR= 1.04 (1.02 to 1.06) - - Diff in % = 12.1 (3.64 to 20.35)MD = 1.50 (1.08 to 1.92) Diff in % = 17.0 (8.83 to 24.94)	None // // // // //

BMI: Body mass index; CLIA: chemiluminescence immunoassay; CLEIA: Electrochemiluminescence immunoassay; CS: Cross-sectional; CV: coefficient of variation; CVD: Cardiovascular disease; Diff in %: percentage difference; DS Polyneuropathy: Distal symmetric polyneuropathy; DKA: Diabetic ketoacidosis; ECLIA: Electrochemiluminescence immunoassay; eGFR: estimated glomerular filtration rate; FBS: Fasting blood sugar; fCP: Fasting C-peptide; Fu.: Follow up; GST: Glucagon stimulation test; HLA: human leukocyte antigens; HR: Hazard ratio; HTN: Hypertension; IA: Immunoassay; MAGE: mean amplitude of glucose excursions; MD: Mean difference; Med: Median; MMTT: mixed meal tolerance test; NA: Not applicable; OR: Odds ratio; PC: Prospective cohort; PCP: Postprandial C-peptide; PP: Postprandial; RC: Retrospective cohort; RR: Relative ratio; rCP: Random C-peptide; RIA: Radioimmunoassay; SD: Standard deviation.

Highlights

- We have summarized the available evidence on the clinical significance of detectable levels of C-peptide in T1DM.
- A systematic search and meta-analysis were performed using online databases.
- Individuals with T1DM in the detectable C-peptide group, compared with the undetectable C-peptide group, had lower mean HbA1c (- 0.08) and daily insulin dose (- 0.41) and showed lower odds for retinopathy (pooled crude odds ratios: 0.53) and nephropathy complications (0.62).
- Individuals with T1DM in the detectable C-peptide group may experience better clinical outcomes.